


ORIGINAL RESEARCH

Journal of Research and Review in Science,

Volume 13, Issue 1, 26-40, June 2026

Doi:



LARVAE BREEDING HABITAT CHARACTERIZATION AND IDENTIFICATION OF *ANOPHELES* MOSQUITOES IN SELECTED LOCAL GOVERNMENT AREAS IN EKITI STATE, NIGERIA

Victor Gboyega¹, Adejumoke Ajayi², Abiodun Adams^{3*}, and Hillary Okoh¹

¹Department of Animal and Environmental Biology, Faculty of Science, Federal University Oye-Ekiti, Ekiti State, Nigeria.

²Department of Integrated Science, School of Science Education, Federal College of Education Ilawe-Ekiti, Ekiti State, Nigeria.

³Department of Zoology and Environmental Biology, Faculty of Science, Lagos State University, Nigeria.

Correspondence

Abiodun Adams,
Department of Zoology and Environmental Biology, Faculty of Science, Lagos State University, Nigeria.
Email: abiodunadamsd@gmail.com

Funding information

The authors received no specific funding for this work. This study was self-funded by the authors.

Abstract

Introduction: Malaria remains a major public health challenge in Nigeria despite ongoing control efforts. Understanding the ecological characteristics of mosquito breeding habitats and the molecular composition of malaria vectors is essential for improving vector control strategies.

Aims: This study aimed to determine the breeding habitat preferences and physicochemical characteristics of *Anopheles* mosquito breeding sites and to identify the molecular composition of members of the *Anopheles gambiae* complex in Ado, Oye, and Ikere LGAs of Ekiti State.

Materials and Methods: A cross-sectional study involving entomological surveys and molecular analyses was conducted. Larval habitats were characterized using WHO standard forms. *Anopheles* larvae were collected from nine breeding sites across the three LGAs and reared to adulthood. Adult mosquitoes were identified morphologically, while sibling species within the *Anopheles gambiae* complex were identified using PCR. Physicochemical parameters (temperature, pH, dissolved oxygen, electrical conductivity, total dissolved solids, and salinity) and essential elements (sodium, calcium, magnesium, copper, iron, phosphate, and nitrogen) were analyzed. Statistical tests were used to compare variations among LGAs.

Results: All breeding habitats were temporary water bodies (100%), with 67% polluted and turbid, and 78% exposed to sunlight. All sites were located near human settlements. Recorded parameters included temperature (25.7–37.8°C), pH (4.0–9.2), dissolved oxygen (16.4–27.4 mg/L), and electrical conductivity (220–554 µS/cm). Significant differences were observed for temperature ($p = 0.02$), pH ($p = 0.03$), and salinity ($p = 0.001$), while EC ($p = 0.39$), TDS ($p = 0.05$), and DO ($p = 0.19$) were not significant. Elemental concentrations varied across habitats. Molecular analysis identified *Anopheles coluzzii* (70.2%), while 29.8% remained unidentified. No significant difference in species distribution was observed ($\chi^2 = 1.696$, $p = 0.428$).

Conclusion: Enhanced vector management strategies, including molecular surveillance and continuous resistance monitoring, are essential for improving insecticide efficacy and malaria control in Ekiti State and other endemic areas.

Keywords: Larvae, *Anopheles* mosquitoes, morphology, polymerase chain reaction, malaria

All co-authors agreed to have their names listed as authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 The Authors. *Journal of Research and Reviews in Science – JRRS*, A Publication of Lagos State University

INTRODUCTION

Malaria is a life-threatening disease caused by Plasmodium parasites, transmitted to humans through the bites of infected Anopheles mosquitoes (WHO, 2023). Despite global efforts to control and eliminate malaria, it remains a significant public health challenge, particularly in sub-Saharan Africa. Nigeria, being one of the countries with the highest malaria burden, has implemented various strategies to combat the disease (WHO, 2022). Ekiti State, located in the south-western region of Nigeria, is not exempt from this burden, with malaria contributing to morbidity and mortality, especially among children under five and pregnant women (Adeyanju *et al.*, 2024). Anopheles mosquitoes, belonging to the genus Anopheles, are the primary vectors responsible for transmitting malaria, a devastating disease caused by Plasmodium parasites (Wondji *et al.*, 2012). There are over 460 recognised species of Anopheles mosquitoes, but only about 30–40 of them are capable of transmitting malaria to humans (WHO, 2016). These mosquitoes exhibit a wide range of biological and ecological characteristics, which significantly influence their capacity to spread malaria (WHO, 2016). The *Anopheles gambiae complex* is one of the most efficient malaria vectors in sub-Saharan Africa (Cisse *et al.*, 2015). In Nigeria, the primary Anopheline mosquito species responsible for malaria transmission include members of the *Anopheles gambiae complex* and the *Anopheles funestus* group. The primary members of this complex are *Anopheles gambiae sensu stricto*, *Anopheles arabiensis*, and *Anopheles coluzzii* (Cisse *et al.*, 2015). *Anopheles gambiae* is highly anthropophilic, preferring to feed on humans, which makes it an effective malaria transmitter (Habibu *et al.*, 2017). *Anopheles gambiae sensu stricto* (s.s.) is one of the most efficient malaria vectors in Nigeria (Akpan *et al.*, 2018). This complex includes several morphologically indistinguishable sibling species, each with different behaviours and ecological adaptations. *Anopheles arabiensis*, on the other hand, exhibits more opportunistic feeding behaviour, feeding on both humans and animals, while *Anopheles coluzzii* is adapted to breeding in permanent water bodies (Habibu *et al.*, 2017). *Anopheles coluzzii*, which was previously considered a molecular form of *Anopheles gambiae* s.s., is now recognised as a distinct species (Ibrahim, 2014). It is adapted to breeding in permanent water bodies and is also an important malaria vector in Nigeria. This research helped collate information; malaria vector control efforts risk becoming ineffective, leading to increased transmission rates and higher disease burden. Understanding the specific Anopheline species present in Nigeria is crucial for designing effective malaria control strategies. Differences in behaviour, breeding habitats, and insecticide resistance patterns among these species require tailored approaches to effectively reduce malaria transmission. For instance, the high anthropophily of *Anopheles gambiae* s.s. necessitates interventions that target human–mosquito contact, such as ITNs, while the varied feeding habits of *Anopheles arabiensis* may require integrated vector management strategies that address both human and animal hosts (Shehu *et al.*, 2023). This study aims to address this gap by investigating mosquito larval habitat characteristics and ascertaining mosquito species present in these areas. This essential data will help inform and optimise local and state-wide malaria vector control programmes.

2.0 MATERIAL AND METHODS

2.1 Study Sites

The study was carried out in three (3) selected local government areas in Ekiti State, located in south-west Nigeria. Ekiti State is situated in the tropical region, bounded by longitudes 40° 51" and 50° 45" East of the Greenwich meridian and latitudes 7° 15" and 8° 51" North of the Equator. The state is divided into 16 local government areas (LGAs), comprising 13 rural, 2 peri-urban and 1 urban area. The state comprises three senatorial districts, namely Ekiti Central, Ekiti South, and Ekiti North. The province covers a total landmass of 6,353 km² and a density of 514 / km². The population of the state is approximately 2.4 million people according to the 2006 population census, but the 2016 estimates put the number of people in the state at 3.3 million (NBS, 2012). The state capital is in Ado-Ekiti, which is both the commercial and political hub of the state (NBS, 2012). The selection of the LGAs was done to include one from each senatorial district and one from urban (Ado LGA), peri-urban (Ikere LGA) and rural (Oye LGA). The study sites were randomly selected within each study location, and their respective coordinates were taken accordingly.




Figure 1: Map of Ekiti state showing the study areas.

2.2 Larval survey

Larvae sampling was conducted biweekly at specified locations from May to June 2024, during which mosquito larvae were collected. At each sampling visit, habitats were initially inspected for the presence of *Anopheles* mosquito larvae, and their coordinates were recorded using a GPS device (Garmin 20.0 eTrex). In addition, the specific locations and classifications of breeding sites were documented for each sampled habitat. Larvae were collected from a variety of sources, including puddles, urban areas, ponds, stagnant water, tree hollows, plant leaf axils, clay pots, discarded vehicle tyres, gutters, tyre tracks, and domestic containers.

2.3 Larvae habitat characterisations

The coordinates of each *Anopheles* mosquito larvae breeding habitat were recorded (Table 1). Habitat characteristics were observed and documented using the WHO characterisation form. Environmental features examined included the habitat's exposure to sunlight, water flow, presence of vegetation and algae, and whether the habitat was natural or artificial, as well as its permanence. Observations also included water clarity and the presence or absence of nearby human settlements. Light intensity was visually classified as either sunlight or shade. The type of substrate was noted, taking into account the presence of vegetation and algae, and whether it was muddy or clear. Water current was visually assessed and categorised as either flowing or stagnant.

Table 1: The study locations, study sites and their respective coordinates

Study Locations	Study Sites	Latitude (N)	Longitude (E)
Ado-Ekiti	Ado poly	7.6111345	5.2807633
	Housing	7.644760	5.251476
	Opopogboro	7.660506	5.228161
Oye-Ekiti	Fallex	7.509404	5.225441
	Egbe	7.508934	5.227866
	Irare	7.530851	5.221124
Ikere-Ekiti	Idi Oshin	7.497687	5.246650
	Faleye	7°34'22.3	5°12'35.9
	Oke Ikere	7°30'01.6	5°13'39.5

2.4 Larvae collection

Habitat characteristics were recorded using a modified WHO larva habitat characterisation form, examining factors such as sunlight exposure, water flow, vegetation, algae presence, and proximity to human settlements. Physicochemical parameters such as temperature, pH, dissolved oxygen (DO), and conductivity were measured on-site with hand-held devices. Standard dipper sampling techniques were employed to collect mosquito larvae from all types of larval habitats, including both man-made and natural sources, at each study location throughout the mosquito breeding season, spanning from May 2024 to June 2024. Sampling was conducted twice a week at each study location. In cases where the use of dippers proved ineffective due to the diminutive size of the habitat, mosquito larvae were collected using Pasteur pipettes. Additionally, the entire water from such habitats was collected using a plastic scoop and transferred into small, clear plastic bowls. To ensure an adequate number of adult mosquitoes for emergence, a strainer with a mesh size of 0.55 mm was used to sieve and consolidate all the collected larvae. This pooling process allowed for a higher concentration of larvae in a single container. Subsequently, the bowls containing the larvae were carefully inspected to identify and remove any unwanted organisms or predators present, employing Pasteur pipettes for precise removal. The collected larvae and pupae were gently placed in clean jam containers, partially filled with water from their respective breeding sites, to ensure a suitable environment for further observation. Each container was meticulously labelled, indicating the specific time, date, and location of collection to facilitate easy identification. The precise locations of the study sites were determined using Global Positioning System (GPS) technology, allowing for accurate coordinate identification.

2.5 Rearing of the Mosquito Larvae to Adult

Mosquito larvae collected were carefully transported in clearly labelled plastic containers to the insectaries at the Entomology Unit of the Department of Animal and Environmental Biology, Federal University Oye-

Ekiti. There, the mosquito larvae were reared to adulthood in the entomology unit. Larvae from different breeding sites were kept separately to ensure accurate species identification. In the laboratory, larvae were placed in individual bowls labelled accordingly and maintained under controlled conditions: room temperature of 27°C and relative humidity of 82%. Adult mosquitoes were fed with yeast, and the water in the bowls was regularly changed to prevent mortality from bacteria or algae growth.

2.6 Sample Preservation

All the *Anopheles* mosquitoes that were subjected to insecticide susceptibility tests were identified morphologically and preserved in labelled Eppendorf tubes containing silica gel desiccant. Samples were kept in the laboratory at a temperature of 27±2°C.

2.7 Morphological Identification of *Anopheles* mosquitoes

Morphological identification was conducted using a stereomicroscope to determine the species group of *Anopheles* mosquitoes. Morphological keys from Gillies and Coetzee (Gillies and Coetzee, 1987) and (Coetzee and Koekemoer, 2013) were utilised as references during the identification process. These keys provided detailed information and characteristics necessary for distinguishing between different species and determining the sex of the mosquitoes accurately. This allows the percentage mortality by species group and species siblings to be properly calculated (WHO, 2013b).

2.8 Molecular Identification of Sibling Species within the *Anopheles gambiae* complex

Molecular identification of sibling species within the *Anopheles gambiae* complex was conducted at the Molecular Entomology and Vector Control Research Laboratory of the Nigerian Institute of Medical Research (NIMR) in Yaba, Lagos State. To identify the sibling species, PCR amplification of DNA extracted from the legs and wings of *Anopheles gambiae* s.s. specimens collected during the study was performed.

2.9 Data analysis

The results were presented as percentages, and the environmental characteristics of habitats were assessed using a chi-square test. The distribution of *Anopheles* species groups, identified through both morphological and molecular methods, was also evaluated using a chi-square test. Differences in knockdown time and other relevant sample collections were assessed using the Wilcoxon Signed-Ranks Test in SPSS version 27. Data were initially entered in Microsoft Excel and subsequently imported into RStudio for analysis. Descriptive statistics, including mean and standard deviation (SD), were calculated to summarise the physicochemical parameters and essential elements of the locations using the group-by function from the dplyr package. Analysis of Variance (ANOVA) was conducted to determine significant differences in means at a 95% confidence level, followed by Tukey's HSD post-hoc test to identify specific group differences. Mean values with similar superscripts across the columns indicate no significant difference.

3.0 RESULTS AND DISCUSSION

3.1 Characterization of larvae breeding habitats for *Anopheles* mosquito across the study locations

Table 2 shows the physio-parameters of waters measured across the sampled sites in the three Local Government Areas (LGAs) of Ado, Oye, and Ikere. The water bodies across Ado-Ekiti, Oye-Ekiti, and Ikere-Ekiti showed distinct characteristics based on their origin, permanency, and environmental features. Water bodies in Ado-Ekiti (Ado Poly, Housing, and Opopogboro) and Ikere-Ekiti (Idi-Osi, Oke, and Faleye) were primarily natural, while Oye-Ekiti (Faleex and Egbe) included man-made sources, except for Irare, which was natural. All the water bodies were temporary, with specific types varying from ponds and ditches to gutters and tyre tracks. Most water currents were stagnant, except for slow-flowing water at Egbe in Oye-

Ekiti. Turbidity varied significantly; Ado-Ekiti had a mix of polluted and clear water, while Ikere-Ekiti showed instances of dark and clear water, and Oye-Ekiti included polluted, clear, and turbid conditions. Substrate types were predominantly muddy across locations, with some clear substrates at Faleye. Settlements were present near all water bodies, and sunlight was the dominant light source, except for Idi-Osi in Ikere-Ekiti, which had no light, and Faleye, which had partial sunlight. Vegetation was present in some areas, particularly in Ado Poly, Housing, Egbe, Oke, and Faleye, while others lacked vegetation. Algae were common in many locations, with exceptions at Egbe, Irare, and Idi-Osi, which showed no algae presence. These variations highlight the diversity in water body characteristics across the study areas.

3.2 Physio-chemical parameters larvae breeding habitats for *Anopheles* mosquito across the study locations

Table 3 presents the physio-chemical parameters of water measured across the sampled sites in the three Local Government Areas (LGAs) of Ado, Oye, and Ikere. For temperature, Ado recorded a mean of $26.2 \pm 1.37^{\circ}\text{C}$, which was significantly lower than Oye's $34.6 \pm 2.94^{\circ}\text{C}$, but overlapped with Ikere's intermediate value of $29.3 \pm 3.26^{\circ}\text{C}$ ($p = 0.02$). The pH in Ado (8.24 ± 0.99) was significantly higher than in Oye (5.07 ± 1.11) and Ikere (6.67 ± 1.16) ($p = 0.03$). Salinity was lowest in Ado (0.35 ± 0.40 ppm), followed by Oye (1.46 ± 0.24 ppm), while Ikere recorded the highest value (2.99 ± 0.72 ppm) ($p = 0.001$). Electrical conductivity (EC) did not vary significantly among locations, with means of 451 ± 137 $\mu\text{S/cm}$ in Ado, 329 ± 45.4 $\mu\text{S/cm}$ in Oye, and 337 ± 129 $\mu\text{S/cm}$ in Ikere ($p = 0.39$). Similarly, total dissolved solids (TDS) measured 350 ± 35.7 mg/l in Ado, 235 ± 37.5 mg/l in Oye, and 388 ± 88.4 mg/l in Ikere, showing partial overlap, with Ikere slightly distinct ($p = 0.05$). Dissolved oxygen (DO) exhibited no significant variation, with mean values of 23.7 ± 3.20 mg/l, 18.9 ± 1.99 mg/l, and 20.1 ± 3.47 mg/l for Ado, Oye, and Ikere, respectively ($p = 0.19$). Overall, temperature ($p = 0.02$), pH ($p = 0.03$), and salinity ($p = 0.001$) showed statistically significant differences among the LGAs, while EC ($p = 0.39$), TDS ($p = 0.05$), and DO ($p = 0.19$) did not.

3.3 Concentration of essential minerals across larvae breeding habitats for *Anopheles* mosquito across the study locations

Table 4 shows that the concentration of essential elements across larvae breeding habitats for *Anopheles* mosquitoes varied significantly among the three LGAs studied. In Ado LGA, sodium (Na) concentrations ranged from 25.2 to 28.9 mg/l, with a mean of 26.6 ± 1.60 mg/l, while calcium (Ca) ranged from 47.4 to 58.3 mg/l, with a mean of 52.4 ± 4.83 mg/l. Magnesium (Mg) levels averaged 10.3 ± 0.81 mg/l, copper (Cu) 0.86 ± 0.06 mg/l, iron (Fe) 0.71 ± 0.16 mg/l, total nitrogen (T.N) 42.3 ± 4.45 mg/l, and total phosphate (T. Phosphate) 10.7 ± 2.19 mg/l. In Oye LGA, sodium concentrations were higher, with a mean of 63.4 ± 8.77 mg/l; calcium averaged 95.9 ± 7.78 mg/l, magnesium 14.3 ± 1.84 mg/l, copper 1.51 ± 0.57 mg/l, iron 2.57 ± 0.54 mg/l, total nitrogen 79.0 ± 19.0 mg/l, and total phosphate 21.5 ± 4.24 mg/l. In Ikere LGA, sodium concentrations ranged widely, with a mean of 28.0 ± 22.3 mg/l; calcium averaged 60.2 ± 28.8 mg/l, magnesium 9.74 ± 3.80 mg/l, copper 0.813 ± 0.24 mg/l, iron 0.74 ± 0.35 mg/l, total nitrogen 45.0 ± 12.5 mg/l, and total phosphate 9.82 ± 1.74 mg/l. Statistical analysis showed significant differences in the mean concentrations of sodium ($P = 0.000$), calcium ($P = 0.001$), magnesium ($P = 0.012$), copper ($P = 0.005$), iron ($P = 0.000$), total nitrogen ($P = 0.001$), and total phosphate ($P = 0.000$) in Oye LGA, compared to the other LGAs.

Table 2: Physical parameters of larvae breeding habitats for *Anopheles* mosquito in the study areas.

	Ado-Ekiti			Oye-Ekiti			Ikere-Ekiti		
Parameters	Ado Poly	Housing	Opopogboro	Faleex	Egbe	Irare	Idi-Osi	Oke	Faleye
Origin of water body	Natural	Natural	Natural	Manmade	Manmade	Natural	Manmade	Natural	Natural
Permanency	Temporary	Temporary	Temporary	Temporary	Temporary	Temporary	Temporary	Temporary	Temporary
Specific water body	Pond	Pond	Ditches	Gutter	Pound	Ditch	Tire track	Pond	Pond
Water current	Stagnant	Stagnant	Stagnant	Stagnant	Slow flowing	Stagnant	Stagnant	Stagnant	Stagnant
Turbidity	Polluted	Polluted	Clear	Polluted	Clear	Turbid	Clear	Polluted	Dark
Substrate type	Mud	Mud	Mud	Mud	Mud	Mud	Mud	Mud	Clear
Presence of settlement	Present	Present	Present	Present	Present	Present	Present	Present	Present
Light intensity	Sunlight	Sunlight	Sunlight	Sunlight	Sunlight	Sunlight	None	Sunlight	Partial
Presence of vegetation	Yes	Yes	No	No	Yes	No	No	Yes	Yes
Algae present	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes

Table 3: Physio-chemical parameters of larvae breeding habitats for *Anopheles* mosquito in the study areas.

LGA	Sampling sites	Temp (°C)	pH	EC (µs/m)	TDS	DO (mg/l)	Salinity (ppm)
Ado	Ado-Poly	27.4	8.3	554	381	27.4	0.25
	Housing	26.5	9.2	295	311	21.8	0.8
	Opopogboro	24.7	7.22	503	358	21.9	0.01
	Mean ± SD	26.2 ± 1.37 ^a	8.24 ± 0.99 ^a	451 ± 137 ^a	350 ± 35.7 ^a	23.7 ± 3.20 ^a	0.35 ± 0.40 ^a
Oye	Faleex	37.8	6.21	319	237	17.7	1.64
	Egbe	34.1	4	290	197	17.8	1.54
	Irare	32	5	379	272	21.2	1.19
	Mean ± SD	34.6 ± 2.94 ^b	5.07 ± 1.11 ^b	329 ± 45.4 ^a	235 ± 37.5 ^a	18.9 ± 1.99 ^a	1.46 ± 0.24 ^{ac}
Ikere	Idi-Oshi	25.7	5.9	475	485	20.5	3.64
	Okekere	32.1	8	315	367	23.3	2.21
	Faleye	30	6.1	220	312	16.4	3.11
	Mean ± SD	29.3 ± 3.26 ^{ab}	6.67 ± 1.16 ^{ab}	337 ± 129 ^a	388 ± 88.4 ^{ab}	20.1 ± 3.47 ^a	2.99 ± 0.72 ^b
	p-value	0.02	0.03	0.39	0.05	0.19	0.001

Table 4: Concentration of essential elements across larvae breeding habitats for *Anopheles* mosquitoes

LGA	Sampling sites	Na (mg/l)	Ca (mg/l)	Mg (mg/l)	Cu(mg/l)	Fe (mg/l)	T.N (mg/l)	T. PHOSPHATE (mg/l)
Ado	Ado Poly (1)	28.3	47.4	11.308	0.782	0.523	44.345	13.516
	Ado Poly (2)	28.9	47.8	11.316	0.778	0.52	44.36	13.509
	Housing (1)	25.8	51.3	9.859	0.876	0.729	36.614	9.274
	Housing (2)	25.4	51.6	9.862	0.882	0.735	36.622	9.278
	Opopogboro (1)	25.2	58.2	9.618	0.912	0.862	45.916	9.249
	Opopogboro (2)	25.8	58.3	9.624	0.917	0.87	45.91	9.253
	Mean $\pm$ SD	26.6 $\pm$ 1.60 ^a	52.4 $\pm$ 4.83 ^a	10.3 $\pm$ 0.81 ^a	0.86 $\pm$ 0.06 ^a	0.71 $\pm$ 0.16 ^a	42.3 $\pm$ 4.45 ^a	10.7 $\pm$ 2.19 ^a
Oye	Faleex	52.8	88.5	12.428	0.926	2.077	55.262	21.307
	Faleex	52.3	88.9	12.433	0.932	2.07	55.256	21.314
	Egbe (1)	66.3	93.2	13.906	1.434	2.387	85.528	16.845
	Egbe (2)	66.2	93.6	13.914	1.428	2.374	85.534	16.842
	Irare (1)	71.6	105.4	16.502	2.178	3.246	96.343	26.311

	Irare (2)	71.5	105.7	16.497	2.19	3.244	96.338	26.316
	Mean $\pm$ SD	63.4 $\pm$ 8.77 ^b	95.9 $\pm$ 7.78 ^b	14.3 $\pm$ 1.84 ^b	1.51 $\pm$ 0.57 ^b	2.57 $\pm$ 0.54 ^b	79.0 $\pm$ 19.0 ^b	21.5 $\pm$ 4.24 ^b
Ikere	Idi-Oshi (1)	10.8	55.8	8.968	0.528	0.624	40.786	10.313
	Idi-Oshi (1)	10.4	55.5	8.972	0.522	0.629	40.792	10.32
	Okekere (1)	16.9	30.3	5.917	0.873	0.415	33.608	7.674
	Okekere (1)	16.5	30.6	5.93	0.879	0.411	33.612	7.681
	Faleye (1)	56.7	94.4	14.315	1.036	1.169	60.59	11.465
	Faleye (1)	56.5	94.3	14.322	1.041	1.16	60.593	11.459
	Mean $\pm$ SD	28.0 $\pm$ 22.3 ^a	60.2 $\pm$ 28.8 ^a	9.74 $\pm$ 3.80 ^a	0.813 $\pm$ 0.24 ^a	0.74 $\pm$ 0.35 ^a	45.0 $\pm$ 12.5 ^a	9.82 $\pm$ 1.74 ^a
	p-value	0.000	0.001	0.012	0.005	0.000	0.001	0.000

3.4 Molecular characterisation of Anopheles species identified across the study LGAs

Table 5 shows that the molecular characterisation of *Anopheles* species across the study LGAs revealed variations in species distribution. In Ado LGA, 16 *Anopheles* mosquitoes were examined, with 11 (68.8%) identified as *An. coluzzii* and 5 (31.2%) remaining unidentified. In Ikere LGA, 15 mosquitoes were examined, of which 9 (60%) were *An. coluzzii* and 6 (40%) were unidentified. Oye LGA had the highest proportion of *An. coluzzii*, with 13 (81.3%) of the 16 mosquitoes examined, leaving 3 (18.7%) unidentified (Figures 1–3). Overall, out of the 47 mosquitoes examined across all LGAs, 33 (70.2%) were identified as *An. coluzzii*, while 14 (29.8%) were unidentified. Statistical analysis showed no significant difference in the distribution of *An. coluzzii* and unidentified species across the LGAs (Chi-square = 1.696, $p = 0.428$). This indicates a relatively consistent pattern of species identification across the study locations.




Figure 2: Electrophoretic gel of molecular identification of members of *An. gambiae* s.l showing the amplification of 463 and 333 bp for *An. Coluzzi* in Ikere LGA

Key: Lane 1 and 19: 100 bp DNA ladder,

Lane 1: 100 bp DNA ladder

Lane 2: Positive control for *Anopheles coluzzii* (463 and 333 bp)

Lane 3: Positive control for *Anopheles gambiae* (463 and 221 bp)

Lane 4: Negative control

Lane 7,8,9,10,14,15,16,17 and 19 are amplified and indicates *Anopheles coluzzii*

Lane 5,6,11,12,13 and 18 were unamplified.

Table 5: Molecular characterization of *Anopheles* species identified across the study LGAs

LGA	Total examined	<i>Anopheles</i>	<i>An. coluzzii</i>	Unidentified	Chi-square	p-value
Ado	16		11 (68.8)	5 (31.2)	1.696	0.428
Ikere	15		9 (60)	6 (40)		
Oye	16		13 (81.3)	3 (18.7)		
Total	47		33 (70.2)	14 (29.8)		

4.0 DISCUSSION

Studies on the habitats of *Anopheles* mosquitoes in Ado, Oye, and Ikere LGAs of Ekiti State showed that temporary water sources are the preferred ones for larval development. This is because the habitats of these water bodies are stagnant, and they are suitable for the ecological requirements of *Anopheles* mosquitoes, as supported by an abundant literature of evidence (Adejanya *et al.*, 2024). Water sources differed considerably, from natural ponds and streams to anthropogenic habitats like gutters, containers that humans had discarded, and tyre tracks. The presence of *Anopheles* larvae in different environments is similar to findings reported by (Ranson and Lissenden, 2016), stressing the need for environmental management to break their breeding cycles. Temperature, pH, and salinity were all found to have important roles in larval development at the three studied LGAs. For example, in Ado, high larval densities were supported by cooler temperatures (26.2°C) and a slightly alkaline pH (8.24). Conversely, Oye's warmer temperatures (34.6°C) and lower pH (5.07) severely curtailed larval growth. Similar correlations between water chemistry and mosquito population dynamics in West Africa have also been reported by (Hernández, 2020), whose findings resonate with those of this study. Polluted habitats around human settlements were also found in previous research to increase the risks of malaria transmission (Ranson and Lissenden, 2016). *Anopheles* larvae develop optimally in organic-rich stagnant water bodies of poor sanitation conditions. The abundance of mineral nutrients and organic pollutants in larval habitats enhances the survival and adaptability of *Anopheles* larvae, particularly in urban environments. Studies have shown that members of the *Anopheles gambiae* complex can thrive in polluted water bodies characterised by high levels of dissolved solids, heavy metals, and organic matter (Awolola *et al.*, 2007; Azrag and Mohammed, 2018). Similar findings have been reported in Nigeria and Senegal, where anthropogenic activities create nutrient-rich breeding sites that promote larval development and complicate malaria control efforts (Sabtiu *et al.*, 2025; Ndiaye *et al.*, 2024). Furthermore, *Anopheles coluzzii* has been consistently identified as the predominant sibling species across multiple African settings, including Nigeria and Ghana (Klinkenberg *et al.*, 2008; Ezeike *et al.*, 2026; Sabtiu *et al.*, 2025). *Anopheles coluzzii* is well adapted to urban and peri-urban habitats and is often predominant in such environments due to its tolerance of polluted and organically enriched breeding sites (Awolola *et al.*, 2007; Omotayo *et al.*, 2022). This species has been reported as the most common member of the *Anopheles gambiae* complex in parts of south-western Nigeria, particularly in urban settings such as Lagos (Omotayo *et al.*, 2022), which aligns with this observation.

4.1 LIMITATIONS OF THE STUDY AND RECOMMENDATIONS FOR FURTHER STUDIES

This paper presents useful insights into the ecological and molecular features of *Anopheles* mosquitoes in selected LGAs of Ekiti State; yet, some limitations should be addressed. The study was limited to three LGAs and a brief sampling period, which may not reflect the complete regional and seasonal variety of mosquito populations and breeding habitats. The extremely restricted sample size for molecular analysis and the failure to identify 29.8% of specimens suggest limitations in the PCR technology utilised and the possible presence of cryptic species. In addition, the study focused only on the *Anopheles gambiae* complex, omitting other significant vectors such as *Anopheles funestus*. Detailed examination of pesticide resistance was also not included, and crucial climatic and socio-ecological issues such as rainfall and land use were not addressed.

Future studies should therefore increase geographical coverage and utilise longitudinal methodologies to capture seasonal fluctuations. The identification of species and the discovery of cryptic diversity will be enhanced by increasing sample size and utilising contemporary molecular tools such as next-generation sequencing. Additional vector species and thorough pesticide resistance profiling should be included in future studies. A more comprehensive understanding of the variables affecting mosquito reproduction and malaria transmission will also result from the integration of environmental and socio-ecological aspects.

4.2 CONCLUSION

The enumeration of mosquitoes across the three Local Government Areas (LGAs) in Ekiti State, South-West Nigeria, facilitated insights into the ecology and molecular composition of Anopheles mosquitoes. Based on the analysis, the following key conclusions were drawn: Anopheles mosquitoes were predominantly found in breeding sites characterised by stagnant water, sunlight exposure, organic material, proximity to settlements, and temporary water sources. Strong correlations were observed between larval density and physio-chemical parameters, supporting their importance in habitat suitability. Morphological identification showed that the *An. gambiae complex* and *An. coluzzii* were the predominant species. Molecular analysis confirmed *An. coluzzii* as the major sibling species across all LGAs. This research contributes important knowledge on malaria vector ecology and species dynamics in Ekiti State, Nigeria, and is critical for malaria control strategies. It determined the main malaria vector species, primarily of the *Anopheles gambiae complex*, such as *An. coluzzii*, and emphasised the need for accurate species identification to target control resources efficiently.

ACKNOWLEDGEMENTS

This study was carried out at the Department of Animal and Environmental Biology, Federal University Oye-Ekiti, Nigeria. We express our gratitude to the reviewers and editors from the Federal College of Education, Ilawe-Ekiti, and Lagos State University for their invaluable suggestions. We also thank Dr. Oyeniyi T.A. and Mrs. Izekor T.R., Laboratory Scientists at the Nigeria Institute of Medical Research (NIMR), for their technical assistance with the molecular aspects of the research. Gratitude is expressed to field and laboratory workers, together with local officials and community members in the designated Local Government Areas of Ekiti State for their collaboration.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Victor Gboyega: Study design, field and lab investigation, original draft, funding acquisition; Adejumo Ajayi: Supported the literature search and writing of the paper; Abiodun Adams: Drafting and Editing Manuscript, data analysis and interpretation; Hillary Okoh: Conceptualization and Supervision, Lab and field supervision. All authors read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

CONSENT

Informed consent was obtained from local community members within the study area, including a designated contact person, prior to the commencement of field activities.

ETHICAL APPROVAL

Research ethics committee approval was obtained from Faculty of Science Ethics committee, Federal University Oye, Ekiti State. In each visit to the study area, an informed consent was gotten from the local people within the vicinity and a contact within the study area was also carried along to show spots with the desired characteristics.

REFERENCES

1. World Health Organization. (2023). *List of WHO prequalified vector control products*. Geneva: World Health Organization. Accessed 29 July 2025. Available: <https://extranet.who.int/pqweb/vector-control-products/prequalified-product-list>
2. World Health Organization. (2022). *World Malaria Report 2022*. Geneva: World Health Organization. Accessed 29 July 2025. Available: <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2022>
3. Adeyanju, A.A., Mbah, N.U., Abass, F.A., & Olutimo, A.L. (2024). Effect of environmental immunity on mathematical modeling of malaria transmission between vector and host population. *Journal of Applied Sciences and Environmental Management*, **28**(1): 205-212. <https://doi.org/10.4314/jasem.v28i1.23>
4. Wondji, C.S., Coleman, M., Kleinschmidt, I., Mzilahowa, T., Irving, H., Ndula, M., Rehman, A., Morgan, J., Barnes, K.G., & Hemingway, J. (2012). Impact of pyrethroid resistance on operational malaria control in Malawi. *Proceedings of the National Academy of Sciences of the United States of America*, **109**(47): 19063–19070. <https://doi.org/10.1073/pnas.1217229109>
5. World Health Organization. (2016). *Test procedures for insecticide resistance monitoring in malaria vector mosquitoes*. World Health Organization. Accessed 29 July 2025. Available: <https://iris.who.int/bitstream/handle/10665/250677/9789241511575-eng.pdf>
6. Cisse, M.B., Keita, C., Dicko, A., Dengela, D., Coleman, J., Lucas, B., ... & Beach, R. (2015). Characterizing the insecticide resistance of *Anopheles gambiae* in Mali. *Malaria journal*, **14**(1): 327. DOI 10.1186/s12936-015-0847-4
7. Habibu, U., Yayo, A.M. and Yusuf, Y.D. (2017). Susceptibility status of *Anopheles gambiae* complex to insecticides commonly used for malaria control in Northern Nigeria. *International Journal of Science and Technology Research*, **6**(6): 47-54.
8. Akpan, G.E., Adepoju, K.A., Oladosu, O.R., & Adelabu, S.A. (2018). Dominant malaria vector species in Nigeria: Modelling potential distribution of *Anopheles gambiae sensu lato* and its siblings with MaxEnt. *PLoS one*, **13**(10): e0204233. <https://doi.org/10.1371/journal.pone.0204233>.
9. Ibrahim, S.S. (2014). High frequency of kdr L1014F is associated with pyrethroid resistance in *Anopheles coluzzi* in Sudan savannah of northern Nigeria. *BMC Infectious Diseases*, **14**(1): 441. <http://www.biomedcentral.com/1471-2334/14/441>
10. Shehu, I.K., Ahmad, H.B., Olayemi, I.K., Solomon, D., Ahmad, A.H., & Salim, H. (2023). Insecticide susceptibility status in two medically important mosquito vectors, *Anopheles gambiae*, and *Culex quinquefasciatus* to three insecticides commonly used in Niger State, Nigeria. *Saudi Journal of biological sciences*, **30**(2): 103-524. <https://doi.org/10.1016/j.sjbs.2022.103524>
11. National Bureau of Statistics (NBS). (2012). *Annual Abstract of Statistics*. Abuja, Nigeria: NBS. Accessed 29 July 2025. Available: <https://www.nigerianstat.gov.ng>
12. Gillies, M.T., & Coetzee, M. (1987). A supplement to the Anophelinae of Africa South of the Sahara. *Publications of the South African Institute for Medical Research*, **55**(Suppl): 1-143.
13. Coetzee, M. and Koekemoer, L. (2013). Molecular systematics and insecticide resistance in the major African malaria vector *Anopheles funestus*. *Annual Review of Entomology*, **58**, 393–412. <https://doi.org/10.1146/annurev-ento-120811-153628>
14. WHO (2013). *Test procedures for insecticide resistance monitoring in malaria vectors mosquitoes*. Geneva, Switzerland: World Health Organization. WHO press; 1-39.
15. Adejayan, A., Okoh, H., & Oyediji, S. (2024). Characterization of Anopheles Mosquitoes Larvae Breeding habitats in Selected Six Local Government Areas in Ekiti State, South-West Nigeria. *ABUAD International Journal of Natural and Applied Sciences*, **4**(1), 18-26. <https://doi.org/10.53982/aijnas.2024.0401.01-j>
16. Ranson, H. and Lissenden, N. (2016). Insecticide resistance in African *Anopheles* mosquitoes: A worsening situation that needs urgent action to maintain malaria control. *Trends in Parasitology*, **32**, 187–196. <http://dx.doi.org/10.1016/j.pt.2015.11.010>

17. Awolola, T.S., Oduola, A.O., Obansa, J.B., Chukwurar, N.J., & Unyimadu, J.P. (2007). *Anopheles gambiae* s.s. breeding in polluted water bodies in urban Lagos, southwestern Nigeria. *Journal of vector borne diseases*, **44**(4): 241–244.
18. Azrag, R.S., & Mohammed, B.H. (2018). *Anopheles arabiensis* in Sudan: a noticeable tolerance to urban polluted larval habitats associated with resistance to Temephos. *Malaria journal*, **17**(1); 204. <https://doi.org/10.1186/s12936-018-2350-1>
19. Sabtiu, A.R.M., Hinne, I.A., Sraku, I.K., Halou, D.K., Doe, R.T., Attah, S.K., ... & Afrane, Y.A. (2025). Larval habitat diversity, physicochemical characteristics and their effect on the larval density of malaria vectors in the city of Accra, Ghana. *Malaria Journal*, **24**(1): 299. <https://doi.org/10.1186/s12936-025-05540-1>
20. Ndiaye, F., Diop, A., Chabi, J., Sturm-Ramirez, K., Senghor, M., Diouf, E.H., ... & Niang, E.H.A. (2024). Distribution and dynamics of *Anopheles gambiae* sl larval habitats in three Senegalese cities with high urban malaria incidence. *Plos one*, **19**(5): e0303473. <https://doi.org/10.1371/journal.pone.0303473>
21. Ezeike, K.A., Danga, S.P.Y., Chukwuma, E.V., Chinaza, F.C., Emmanuel, O.O., Chukwuebuka, M.N., ... & Esimone, C.O. (2026). Urban evolution of insecticide resistance and susceptibility patterns of *Anopheles coluzzii* in Southeastern Nigeria: implications for malaria vector control. *Malaria Journal*. **25**(86): 12. <https://doi.org/10.1186/s12936-025-05760-5>
22. Klinkenberg, E., McCall, P.J., Wilson, M.D., Amerasinghe, F.P., & Donnelly, M.J. (2008). Impact of urban agriculture on malaria vectors in Accra, Ghana. *Malaria journal*, **7**(1): 151. [doi:10.1186/1475-2875-7-151](https://doi.org/10.1186/1475-2875-7-151)
23. Omotayo, A.I., Ande, A.T., Oduola, A.O., Adelaja, O.J., Adesalu, O., Jimoh, T.R., ... & Awolola, S.T. (2022). Multiple insecticide resistance mechanisms in urban population of *Anopheles coluzzii* (Diptera: Culicidae) from Lagos, South-West Nigeria. *Acta Tropica*, **227**, 106291. <https://doi.org/10.1016/j.actatropica.2021.106291>